


Reduction of a benzonitrile; 1-[4-(trans-4-pentylcyclohexyl)phenyl]methanamine

SyntheticPage 708

Submitted Dec 16, 2013, published Dec 17, 2013

DOI: [10.1039/SP708](https://doi.org/10.1039/SP708)

John MacMillan (john.macmillan@temple.edu)



Chemicals Used

4-(Trans-4-pentylcyclohexyl)-benzonitrile (Fluorochem), 300840

Lithium aluminum hydride, Sigma Aldrich, powder, 95%, 19,987-7. freshly opened can

Ether, Sigma Aldrich, anhydrous, 99.8%, 29,608-2, freshly opened bottle

Sodium sulfate, Sigma Aldrich, anhydrous, 99+%, 23,931-3

Procedure

An oven dried 100ml three neck flask, equipped with magnetic stirrer, water condenser and nitrogen inlet was charged under nitrogen with 1.0 g (4 mmol) 4-(trans-4-pentylcyclohexyl)-benzonitrile (Fluorochem) and 50 ml of anhydrous ether. The mixture was stirred until all of the nitrile had dissolved. 1.0 g of freshly opened powdered lithium aluminum hydride was then added through a powder funnel and the mixture stirred overnight. A 10ml pressure equalized dropping funnel was then attached and the excess hydride decomposed by drop wise addition of 5 ml water over one hour. A copious white precipitate of aluminum hydroxide was then present. The precipitate was filtered off with a small Buchner funnel under water aspirator pressure and washed under vacuum with 5 ml ether. The filtrate of ether/amine was dried for one hour over 2 grams of anhydrous sodium sulfate and the ether solution decanted into a 100 ml round bottom flask. The sulfate residue was twice stirred, utilizing a glass rod, with two 5 ml ether portions which were each decanted and combined with the filtrate. Ether then was removed on a rotovap (water bath ~80⁰C) and the residue pipetted into a 2 ml conical distilling flask equipped with a short path distilling head and an oil bath. Distillation under full rotary pump vacuum (bath temperature ~150⁰), yielded 800 mg (80%) of the liquid crystalline amine, which solidified on standing at room temperature. Yields of all liquid crystalline amines in these references are in the 70-85% range, with losses mostly mechanical.

Author's Comments

Caution! Lithium aluminum hydride is extremely water sensitive. Use a freshly opened can if possible and weigh in glovebox if possible. On contact with excess water it decomposes violently with hydrogen evolution. It should always be decomposed in a reaction mixture when necessary by slow dropwise addition of water. This and other amines described in the primary and secondary references are hygroscopic and should be stored in desiccators. The methanamines may be converted to N-alkyl substituted liquid crystalline cyclohexylphenyl amines or biphenyl amines by further reaction with acid chlorides and lithium aluminum hydride reduction (see references).

Data

B.p. 135⁰(0.1 mm),

m.p. crystal-isotropic (K-I), 27⁰, cooling, isotropic-smectic-A, (I-S_A) , 9⁰, Smectic-A-crystal, (S_A-K), 0⁰.

I.r. (cm-1) 3300,3250,3010,2900,1600

Analysis: Calculated for C₁₈H₂₉N: C, 83.32, H, 11.27, N, 5.40. Found: C, 83.22, H, 11.40, N, 5.46

Lead Reference

John H. MacMillan and Mortimer M. Labes, "Low Transition Temperature Liquid Crystalline Amines Incorporating the Trans-1,4-Cyclohexane Ring System", Molecular Crystals and Liquid Crystals, Vol. 55, p 61, (1979). DOI: [dx.doi.org/10.1080/00268947908069791](https://doi.org/10.1080/00268947908069791)

Other References

John H. MacMillan and Mortimer M. Labes, "Low Transition Temperature Liquid Crystalline Amines Incorporating the Biphenyl Ring System", Mol. Crystals and Liquid Crystals Letters, Vol. 56, p51, (1979). DOI: Link: <http://dx.doi.org/10.1080/01406567908071966>

Chemspider deposition: <http://www.chemspider.com/Chemical-Structure.18536103.html>

John H. MacMillan and Mortimer M. Labes, "Amine Substituted Liquid Crystal Compositions", U.S. Patent 4,293,193, Oct. 6, 1981.

Keywords: amines, hydrogenation